



## Health, Healthy Living and Seniors

Public Health and Primary Health Care  
Communicable Disease Control  
4th Floor, 300 Carlton St, Winnipeg, MB R3B 3M9  
T 204 788-6737 F 204 948-2040  
[www.manitoba.ca](http://www.manitoba.ca)

November, 2015

### Re: Cryptosporidiosis Reporting and Case Investigation

Reporting of cryptosporidiosis (*Cryptosporidium* species) is as follows:

#### Laboratory:

- All positive laboratory results for *Cryptosporidium* species are reportable to the Public Health Surveillance Unit by secure fax (204-948-3044).

#### Health Care Professional:

- Probable (clinical) cases of cryptosporidiosis are reportable to the Public Health Surveillance Unit using the *Clinical Notification of Reportable Diseases and Conditions* form (<http://www.gov.mb.ca/health/publichealth/cdc/protocol/form13.pdf>) ONLY if a positive lab result is not anticipated (e.g., poor or no specimen taken, person has recovered).
- Cooperation in Public Health investigation is appreciated.

#### Regional Public Health or First Nations Inuit Health Branch (FNIHB):

- Once the case has been referred to Regional Public Health or FNIHB, the *Communicable Disease Control Investigation Form* ([www.gov.mb.ca/health/publichealth/cdc/protocol/form2.pdf](http://www.gov.mb.ca/health/publichealth/cdc/protocol/form2.pdf)) should be completed and returned to the Public Health Surveillance Unit by secure fax (204-948-3044).

Sincerely,

“Original Signed By”

Richard Baydack, PhD  
Director, Communicable Disease Control  
Public Health and Primary Health Care  
Manitoba Health, Healthy Living and Seniors

“Original Signed By”

Carla Ens, PhD  
Director, Epidemiology & Surveillance  
Public Health and Primary Health Care  
Manitoba Health, Healthy Living and Seniors

## Cryptosporidiosis



Public Health Branch

### 1. Case Definition

#### 1.1 Confirmed Case

Laboratory confirmation of infection from an appropriate clinical specimen (e.g., stool, intestinal fluid or small bowel biopsy) with or without clinical illness<sup>a</sup>:

- Demonstration of *Cryptosporidium* oocysts  
OR
- Detection of *Cryptosporidium* DNA  
OR
- Demonstration of *Cryptosporidium* antigen by an approved method (e.g., enzyme immunoassay) (1).

#### 1.2 Probable Case

Clinical illness<sup>a</sup> in a person who is epidemiologically linked to a confirmed case (1).

### 2. Reporting and Other Requirements

#### Laboratory:

- All positive laboratory results are reportable to the Public Health Surveillance Unit (204-948-3044 secure fax).
- Clinical laboratories are required to submit only requested residual specimens from individuals who tested positive for *Cryptosporidium* to Cadham Provincial Laboratory (CPL) within seven days of request.

#### Health Care Professional:

- Probable cases are reportable to the Public Health Surveillance Unit (form available at: [www.gov.mb.ca/health/publichealth/cdc/protocol/form2.pdf](http://www.gov.mb.ca/health/publichealth/cdc/protocol/form2.pdf)) ONLY if a positive lab result is not anticipated (e.g., poor or no specimen taken, person has recovered). Confirmed cases do not require reporting by health care professional as they will be reported to Manitoba Health by the laboratory.

### 3. Clinical Presentation

The major symptom is diarrhea, which may be profuse and watery, accompanied by cramping abdominal pain (2). General malaise, fever, anorexia, nausea and vomiting occur less often; however, diarrhea may be preceded by anorexia and vomiting in children (2). In immunocompetent individuals, symptoms may be intermittent but usually resolve within 30 days (2). The duration of illness in most cases is approximately five to 10 days (3). Asymptomatic infections are common and can be a source of infection for others (2). Disease severity depends on the host (concurrent infection, malnutrition, immunosuppression), agent (genotypic strain), and environmental (dose of exposure) factors (4). Immunocompromised individuals may experience prolonged severe diarrhea which may lead to malnutrition and weight loss (5). Disease can be life-threatening in some immunocompromised hosts (6). Clinical illness is more frequent and severe among the elderly (3). The main site of infection is the small intestine, although infection may be spread throughout the gastrointestinal tract and extra-intestinal sites (7).

Evidence is emerging that the clinical presentation may vary with infecting species (7). There is some evidence that illness is more severe in patients infected with *C. hominis* than those infected with *C. parvum* (8). Recurrence of intestinal symptoms after resolution of the acute stage of illness (for both *C. parvum* and *hominis*) may occur (9). *C. hominis* is associated with increased risk of non-intestinal postinfectious symptoms (e.g., joint pain) (9).

---

a Characterized by diarrhea (often profuse and watery), abdominal cramps, anorexia, fever, nausea, general malaise and vomiting.

## 4. Etiology

*Cryptosporidium parvum* and *C. hominis*, oocyst-forming, intracellular coccidian protozoa, are the two species most often associated with human infection (2, 5, 10-12). There are at least 16 established *Cryptosporidium* species (8).

## 5. Epidemiology

### 5.1 Reservoir

Humans, cattle and other domesticated and feral animals (2). *Cryptosporidium* does not multiply in foods; however, it can retain viability, and therefore infectivity, under moist and cool conditions for several months (13).

### 5.2 Transmission

Transmission is fecal-oral (2). Fecal-oral transmission can occur directly through person-to-person and animal-to-person routes or indirectly through environmental vehicles including water and food (7, 14, 15). The infectious dose is low, facilitating transmission from sources with low-grade contamination, such as recreational water (3). Oocysts can survive adverse environmental conditions for long periods of time (2) and are resistant to commonly used disinfectants (e.g., chlorine and chlorine derivatives) (2, 3, 16, 17). Outbreaks have been associated with day care centres, municipal drinking water and recreational water venues (e.g., swimming pools, waterslides, lakes) (2, 14, 15, 18-20), travel, and contact with farm animals (12, 15, 21). Outbreaks have also been reported in health care facilities (15). Transmission is associated with anal-genital sex (3). At least one study has suggested that sexual behaviour in men who have sex with men is a significant risk factor for cryptosporidial diarrhea (22).

### 5.3 Occurrence

**General:** Worldwide. In industrialized countries, prevalence of infection is less than 1% to 4.5% based on stool examination (2). Prevalence ranges from 3% to 20% in developing regions (2).

Infection is more common in children under two years of age (2, 7), animal handlers, travelers, men who have sex with men (MSM) and close personal contacts of infected individuals (families, health care and day care workers) (2). There is also an increased frequency of infection in patients with altered cellular immunity (3). Surveillance studies have indicated that males were more frequently infected than females; this could be related in part to the over-representation of HIV-infected patients who in at least one study were mainly male (23).

**Canada:** In 2010, 136 cases of *Cryptosporidium* were reported to the National Enteric Surveillance Program (NESP) (24). Of the cases reported, one was identified as being travel-associated (24). *Cryptosporidium* infections are not routinely reported to the provincial or central reference laboratories and are greatly under-represented in NESP (24).

**Manitoba:** In 2011, 19 cases of cryptosporidiosis were reported to Manitoba Health. For 2000-2011 inclusive, 536 cases were reported with the annual incidence rate varying from 1.5 to 8.6 per 100,000 population. For 2000-2011, the highest number of cases (204) was reported in the 5-19 year age group.

### 5.4 Incubation Period

The average incubation period is seven days, but ranges from one to 12 days (2).

### 5.6 Susceptibility and Resistance

Immunocompetent individuals may have asymptomatic infections or self-limited symptomatic infections (2). Adults and children can have multiple episodes of cryptosporidiosis, indicating that acquired immunity to *Cryptosporidium* infection is short-lived or incomplete (7, 11). Children in developing countries are vulnerable to persistent infection because of the independent and synergistic effects of immune naïveté, malnutrition, and HIV infection (25).

## 5.7 Period of Communicability

Oocysts are excreted in stool from the onset of symptoms until several weeks after resolution of symptoms and remain infectious in the environment for two to six months or longer if the environment is moist (2). In immunocompetent people, oocyst shedding usually ceases within two weeks of symptom resolution (2). In immunocompromised people, the period of oocyst shedding can continue for months (5).

## 6. Laboratory Diagnosis

Three stool specimens should be collected on three separate days at least 24 hours apart, but within a 7-10 day time interval. Diagnosis is generally made by identification of oocysts in fecal smears or intestinal biopsy sections. Because shedding can be intermittent, at least three stool specimens should be examined before considering test results to be negative.

New and more sensitive enzyme immunoassay (EIA)-based methods for direct antigen detection in stool have recently become available, but are not routinely available in Manitoba.

## 7. Key Investigations for Public Health Response

- History
  - Child care facility contact;
  - Recreational water contact;
  - Animal contact including farm or companion animals, petting zoos;
  - Similar illness in household or other close contact; and
  - Travel history.
- Determine if there is a common vehicle responsible for other cases such as contaminated drinking water or food. The medical lead for contaminated drinking water issues is the regional Medical Officer of Health. Water sampling would be undertaken by the Office of Drinking Water with assistance from public health inspectors. Public Health Inspectors are

responsible for the investigation of contaminated recreational water. The Public Health Inspector recreational water specialist can be contacted at 204-945-0835.

- Microscopic examination of feces in symptomatic contacts (2).

## 8. Control

### 8.1 Case Management

#### Treatment:

- Specific therapy is generally not indicated in immunocompetent individuals (5).
- Fluid and electrolyte replacement with oral rehydration solutions or intravenous fluids may be required (2).
- If the patient is taking immunosuppressive drugs, consultation with the physician treating the immunosuppression is recommended to determine if these drugs can be stopped or the dose reduced while the patient is ill with cryptosporidiosis (2, 26).
- In patients co-infected with human immunodeficiency virus (HIV), improvement in CD4+ T-lymphocyte levels associated with antiretroviral therapy can lead to symptom resolution and cessation of oocyst shedding (5, 6).
- There is some evidence that a three day course of nitazoxanide treatment reduces the duration of both diarrhea and oocyst shedding in immunocompetent individuals (27-29). In Canada, nitazoxanide is only available through Health Canada's Special Access Program at <http://www.hc-sc.gc.ca/dhp-mps/acces/drugs-droques/index-eng.php>.

#### Infection Control Measures:

- Refer to page 84 of the Manitoba Health document *Routine Practices and Additional Precautions: Preventing the Transmission of Infection in Health Care* available at: <http://www.gov.mb.ca/health/publichealth/cdc/docs/ipc/rpap.pdf>.
- *Cryptosporidium* is not killed by alcohol gels and hand sanitizers (30).

## Public Health Measures:

- Infected individuals should wash their hands frequently for a minimum of 20 seconds using friction with **soap and running water**, especially before preparing food or after going to the toilet, and avoid close contact with anyone with a weakened immune system (4).
- Exclusion of symptomatic individuals from food handling and from direct care of hospitalized and institutionalized patients until asymptomatic (2). Stress proper hand washing (2).
- Exclusion of symptomatic children and child care facility workers who work directly with children from child care settings until diarrhea has resolved (2, 5).
- It is recommended that infected persons do not use recreational water venues (e.g., swimming pools, wading pools, whirlpools) until two weeks after resolution of symptoms (5, 20, 31).
- If waterborne transmission is suspected, large volume water sampling filters can be used to look for oocysts in the water (2).

## 8.2 Contact Management

- Microscopic examination of feces in symptomatic contacts (2).
- Education on preventive measures should occur (refer to Section 8.4 below).

## 8.3 Outbreak Management

An outbreak is defined as the occurrence of case(s) in a particular area and period of time in excess of the expected number of cases.

- Outbreaks should be investigated to identify a common source of infection and prevent further exposure to that source. The extent of the investigation will depend upon the number of cases, the likely source of contamination, and other factors.
- Refer to the *Enteric Illness Protocol* available at: [www.gov.mb.ca/health/publichealth/cdc/protocol/enteric.pdf](http://www.gov.mb.ca/health/publichealth/cdc/protocol/enteric.pdf)
- Public notification may be appropriate and will depend on the number of cases, the likely source of contamination and other factors. The level of

notification will usually be at the discretion of regional Public Health and/or provincial Public Health for local outbreaks but may be at the discretion of the Federal Government for nationally linked foodborne outbreaks as per *Canada's Foodborne Illness Outbreak Response Protocol (FIORP) 2010: To guide a multijurisdictional response* available at: [www.phac-aspc.gc.ca/zoono/fiorp-pritioa/index-eng.php](http://www.phac-aspc.gc.ca/zoono/fiorp-pritioa/index-eng.php)

- *Cryptosporidium* is not killed by alcohol gels and hand sanitizers so these materials are of little use in controlling an outbreak (30).
- Education on preventive measures should occur (refer to Section 8.4 below).

## 8.4 Preventive Measures

- Public education in personal hygiene. All individuals should wash hands well for a minimum of 20 seconds using friction with **soap and running water**:
  - Before preparing, serving, or eating food;
  - After using or cleaning the toilet or assisting someone else with using the toilet;
  - After changing a diaper;
  - After caring for someone who is ill with diarrhea, and
  - After handling an animal or its waste (5).
- Exclusion of symptomatic individuals from food handling and from direct care of hospitalized and institutionalized patients until asymptomatic (2).
- Exclusion of symptomatic children and child care facility workers who work directly with children from child care settings until diarrhea has resolved (2, 5).
- Adoption by water utilities of improved disinfection technologies that inactivate *Cryptosporidium* (14).
- Documentation and remediation following fecal accidents in pools is an essential component in a strategy to reduce the spread of recreational water illnesses (20).



- Drinking water can be decontaminated by boiling [the time of boiling is dependent on altitude (one minute at sea level)] (5) or by filtration with a  $\leq 1\mu\text{m}$  pore size (3).

## References

1. Public Health Agency of Canada. Case Definitions for Communicable Diseases under National Surveillance. *Canada Communicable Disease Report CCDR* 2009; 35S2: 1-123.
2. Heymann David L. Cryptosporidiosis. In: *Control of Communicable Diseases Manual 19th ed*, American Public Health Association, Washington, 2008; 157-159.
3. Petri Jr. WA and Haque R. *Cryptosporidium* Species. In: Mandell GL, Bennett JE, Dolin R eds. *Principles and Practice of Infectious Diseases 7th ed*. Elsevier, Philadelphia, 2010.
4. Weir E. The cryptic nature of cryptosporidiosis. *Canadian Medical Association Journal* 2001; 164(12): 1743.
5. American Academy of Pediatrics. Cryptosporidiosis. In: Pickering LK ed. *Redbook 2012 Report of the Committee on Infectious Diseases 29th ed*. Elk Grove Village, IL: American Academy of Pediatrics, 2012; 296-298.
6. Clark DP. New Insights into Human Cryptosporidiosis. *Clinical Microbiological Reviews* 1999; 12(4): 554-563.
7. Chalmers RM and Davies AP. Minireview: Clinical cryptosporidiosis. *Experimental Parasitology* 2010; 124: 138-146.
8. Cama VA, Bern C, Roberts J *et al*. *Cryptosporidium* Species and Subtypes and Clinical Manifestations in Children, Peru. *Emerging Infectious Diseases* 2008; 14(10): 1567-1574.
9. Hunter PR, Hughes S, Woodhouse S *et al*. Health Sequelae of Human Cryptosporidiosis in Immunocompetent Patient. *Clinical Infectious Diseases* 2004; 39: 504-510.
10. Xiao L. Molecular epidemiology of cryptosporidiosis: An update. *Experimental Parasitology* 2010; 124: 80-89.
11. Xiao L, Fayer R, Ryan U and Upton SJ. *Cryptosporidium* Taxonomy: Recent Advances and Implications for Public Health. *Clin. Microbiol. Rev.* 2004; 17(1): 72-97.
12. Hunter PR, Hadfield SJ, Wilkinson D *et al*. Subtypes of *Cryptosporidium parvum* in Humans and Disease Risk. *Emerging Infectious Diseases* 2007; 13(1): 82-88.
13. Sunnote O, Lowery CJ, Moore JE *et al*. *Cryptosporidium*. *Letters in Applied Microbiology* 2006; 43: 7-16.
14. Centers for Disease Control and Prevention. Cryptosporidiosis Outbreaks Associated with Recreational Water Use – Five States, 2006. *Morbidity and Mortality Weekly Report MMWR* 2007; 56(29): 729-732.
15. Semenza JC and Nichols G. Cryptosporidiosis Surveillance and Water-Borne Outbreaks in Europe. *Eurosurveillance* 2007; 12: 120-123.
16. Naumova EN, Egorov AI, Morris RD and Griffiths JK. The Elderly and Waterborne *Cryptosporidium* Infection: Gastroenteritis Hospitalizations before and during the 1993 Milwaukee Outbreak. *Emerging Infectious Diseases* 2003; 9(4): 418-425.
17. Health Canada. North Battleford, Saskatchewan Spring, 2001. Waterborne Cryptosporidiosis Outbreak, September 24, 2001. Available at: <http://www.health.gov.sk.ca/nb-waterborne-cryptosporidiosis-outbreak>
18. Shields JM, Hill VR, Arrowood MJ and Beach MJ. Inactivation of *Cryptosporidium parvum* under chlorinated recreational water conditions. *Journal of Water and Health* 2008; 513-520.
19. Public Health Agency of Canada. Waterborne Cryptosporidiosis Outbreak, North Battleford, Saskatchewan, Spring 2001. *Canada Communicable Disease Report CCDR* 2001; 27(22): 185-192.

20. Public Health Agency of Canada. An Outbreak of *Cryptosporidium parvum* in a Surrey Pool with Detection in Pool Water Sampling. *Canada Communicable Disease Report CCDR* 2004; 30 (7): 61-66.
21. Chalmers RM and Giles M. Zoonotic cryptosporidiosis in the UK – challenges for control. *Journal of Applied Microbiology* 2010; 109: 1487-1497.
22. Hellard M, Hocking J, Willis J *et al.* Risk factors leading to *Cryptosporidium* infection in men who have sex with men. *Sex Transm Infect* 2003; 79: 412-414.
23. The ANOFEL *Cryptosporidium* National Network. Laboratory-based surveillance for *Cryptosporidium* in France, 2006-2009. *Eurosurveillance* 2010; 15(33):pii=19642.
24. Public Health Agency of Canada. National Enteric Surveillance Program (NESP) Annual Summary 2010. Available at:  
[http://www.nml-hnm.gc.ca/NESP-PNSME/assets/pdf/NESP\\_2010\\_Annual\\_Report\\_ENG.pdf](http://www.nml-hnm.gc.ca/NESP-PNSME/assets/pdf/NESP_2010_Annual_Report_ENG.pdf)
25. Mor SM and Tzipori S. Cryptosporidiosis in Children in Sub-Saharan Africa: A Lingering Challenge. *Clinical Infectious Diseases* 2008; 47: 915-921.
26. Centers for Disease Control and Prevention. *Cryptosporidium* Infection – Immunocompromised Persons. Available at:  
[http://www.cdc.gov/parasites/crypto/gen\\_info/infect\\_ic.html](http://www.cdc.gov/parasites/crypto/gen_info/infect_ic.html) .
27. Rossignol JA, Ayoub A and Ayers MS. Treatment of Diarrhea Caused by *Cryptosporidium parvum*: A Prospective Randomized, Double-Blind, Placebo-Controlled Study of Nitazoxanide. *The Journal of Infectious Diseases* 2001; 184: 103-106.
28. Rossignol J, Kabil SM, El-Gohary Y and Younis AM. Effect of Nitazoxanide in Diarrhea and Enteritis Caused by *Cryptosporidium* Species. *Clinical Gastroenterology and Hepatology* 2006; 4: 320-324.
29. Abubakar II, Aliyu SH, Arumugan C *et al.* Prevention and treatment of cryptosporidiosis in immunocompromised patients (Review). *The Cochrane Library* 2012, Issue 5.
30. Centers for Disease Control and Prevention. Intensified Cryptosporidiosis (Crypto) Control Measures for the Child Care Setting. Available at:  
<http://www.cdc.gov/parasites/crypto/daycare/outbreak.html> .
31. Centers for Disease Control and Prevention. Nitazoxanide: Is there a treatment for diarrhea caused by *Cryptosporidium*? Available at:  
[http://www.cdc.gov/parasites/crypto/health\\_professionals/tx.html](http://www.cdc.gov/parasites/crypto/health_professionals/tx.html) .