

RSV Vaccine Quick Reference Guide for Immunizers

RSV	RSVPreF3 Subunit Vaccine	ABRYSSVO™ Pfizer: 60 years of age and older AND living in a Personal Care Home		
Product Monograph: b3cae670-abaf-4567-94de-6c5997272d2b (pfizer.com) Fact Sheet: www.gov.mb.ca/health/publichealth/factsheets/rsv.pdf Manitoba Health's RSV Webpage: www.gov.mb.ca/health/publichealth/diseases/rsv.html Eligibility Criteria: www.gov.mb.ca/health/publichealth/cdc/vaccineeligibility.html Canadian Immunization Guide: www.canada.ca/en/public-health/services/publications/healthy-living/canadian-immunization-guide-part-4-active-vaccines/respiratory-syncytial-virus.html				
Presentation	Storage & Handling	Administration	Potential Allergens	
<p>Supplied: 0.5ml single dose vial Requires dilution with 0.65ml sterile water (supplied)</p> <p>Abrysvo™ is supplied as a carton containing 10 vials of powder, 10 pre-filled syringes of diluent and 10 vial adapters.</p> <p>The vaccine is a lyophilized white powder for solution provided in a single-dose vial that must be reconstituted with sterile water diluent before use. The final reconstituted solution is clear and colourless. Completely and slowly withdraw the entire contents into the syringe to ensure a 0.5 mL dose of Abrysvo™.</p>	<p>Storage: 2° to 8°C Do not freeze <i>Discard product if exposed to freezing</i></p> <p>Protect from light</p> <p>Use as soon as possible after removal from refrigeration</p> <p><u>After reconstitution:</u> Abrysvo™ should be administered immediately (within 4 hours) after reconstitution. Store the reconstituted vaccine between 15°C and 30°C. Do not freeze reconstituted vaccine.</p> <p>Do not use this vaccine after the expiry date (the expiry date refers to the last day of that month).</p>	<p>Dose: 0.5ml dose</p> <p>Route: Intramuscular (IM) Deltoid</p> <p>Schedule: 1 dose <i>If previously immunized with a RSV vaccine, an additional dose is not recommended.</i></p> <p><i>It is recommended that personal care homes plan to administer the RSV vaccine minimum of two (2) weeks after the administration of influenza vaccines. However, the RSV vaccine may be administered at the same time as other vaccines on an individual resident basis.</i></p> <p>Recommended Needle Gauge and Length: 22-25G 1"- 1 ½"</p>	<p>Mannitol, polysorbate 80, sodium chloride, sucrose, tromethamine, trometamol hydrochloride</p> <p>LATEX FREE</p>	