

Influenza Surveillance 2016–2017

Week 52 & 01 (Dec.25, 2016–Jan.7, 2017)

Data extracted Jan.13, 2017 at 11:00 am

Next report date: Jan. 20, 2017

Laboratory-confirmed influenza A cases: ↑

- Week 52: 2
- Week 01: 13

Laboratory-confirmed influenza B cases: ↑

- Week 52: 4
- Week 01: 1

Since Sept. 1, 2016:

- Influenza A cases: 38
- Influenza B cases: 7

Laboratory

Cases and incidence rates (cases per 10,000 population) since Sept. 1, 2016:

- Winnipeg: 24 (3.2)
- Southern: 4 (2.1)
- Interlake-Eastern: 3 (2.4)
- Prairie Mountain: 14 (8.3)
- Northern: 0 (0.0)

* Rates were not calculated due to small numbers.

Regional Health Authority

Severe outcomes associated with laboratory-confirmed diagnosis of influenza in Week 52 and 01: ↑

- Hospitalizations: 5
- ICU admissions: 1
- Deaths: 0

Since Sept. 1, 2016:

- Hospitalizations: 9
- ICU admissions: 3
- Deaths: 1

*ICU admissions were also included in hospitalizations.

Severity

Calls to Influenza Service at Health Links–Info Santé:

- Week 52: 10
- Week 01: 11

Percent of visits to sentinel physicians due to ILI: ↑

- Week 52: 1.7%
- Week 01: 1.5%

In Emergency Department, ILI patients per day: ↑

- Week 52: 191
- Week 01: 175

Influenza-Like-Illness (ILI)

Laboratory-confirmed Influenza A outbreaks: ↑

- Week 52: 0
- Week 01: 2

Laboratory-confirmed Influenza B outbreaks:

- Week 52: 0
- Week 01: 0

Since Sept. 1, 2016:

- Influenza A outbreaks: 3
- Influenza B outbreaks: 0

Outbreak

Units of Oseltamivir dispensed from community retail pharmacies: ↑

- Week 52: 43
- Week 01: 58
- Since Oct. 1, 2016: 497

Isolates resistant to antiviral since Sept. 1, 2016 in Manitoba:

- Oseltamivir: 0
- Zanamivir: 0

Antiviral

As of Jan. 7, 2017:

- Percent of Manitoban residents immunized with the seasonal influenza vaccine: 19.5%

Immunization

The seasonal influenza activity level in Manitoba overall is still low for this time of year but it is increasing. Two influenza outbreaks in long term care facilities and five influenza associated hospital admissions were reported after Christmas. Older people are affected more than other age groups this season. Over half of the influenza patients are over the age of 65. ILI activity in the province has also been increased. However, respiratory syncytial virus (RSV) continues to be the most commonly detected virus, accounting for over 70% of all respiratory detections last week.

Nationally, seasonal influenza activity continues to increase with more influenza detections, hospital admissions and outbreaks. Some provinces including British Columbia, Alberta, and Ontario have reported high influenza activity.

Globally, influenza A(H3N2) virus has been predominant. The majority of influenza viruses characterized so far are antigenically similar to the vaccine component for 2016-2017. In the United States, ILI activity just surpassed the seasonal threshold.

Summary: Increasing activity

Note. Numbers are subject to change. Missed events in the current report due to a delay of submission to MHSAL will be included in later reports when data become available.

Laboratory Surveillance

Reports of influenza nucleic acid detection, culture isolation, and enzyme immunoassay (EIA) detections are received from Cadham Provincial Laboratory (CPL) and occasionally other laboratories. These reports are forwarded to Epidemiology and Surveillance (E&S) within 24 hours of confirmation. CPL also performs testing for other respiratory viruses including parainfluenza, RSV, adenovirus, rhinovirus, coronavirus, enterovirus, and bocavirus, which are reported to E&S on a weekly basis.




Figure 1. Weekly incidence of lab-confirmed influenza A, Manitoba, 2016–2017




Figure 2. Weekly incidence of lab-confirmed influenza B, Manitoba, 2016–2017




Figure 3. Influenza A and B cases by age group, Manitoba, 2016–2017

Outbreak

Outbreaks are directed to E&S by a phone call or email from public health staff within Regional Health Authorities (RHAs) or from CPL advising the assignment of an outbreak code. CPL submits both positive and negative laboratory results related to outbreaks to E&S. Outbreak investigations are reported from RHAs to E&S by completing an outbreak summary report form on paper or electronically through the Canadian Network for Public Health Intelligence (CNPHI).




Figure 4. Weekly lab-confirmed influenza and ILI outbreaks, Manitoba, 2016-2017

Health Links – Info Santé

Health Links–Info Santé is a 24-hour, 7-days a week telephone information service. It is staffed by registered nurses with the knowledge to provide answers to health care questions and guidance to appropriate care over the phone. When a caller phones Health Links–Info Santé and selects Influenza Service, they are given an option to select information on (1) the groups of individuals who are at an increased risk of serious illness, (2) how to arrange an influenza vaccine, (3) the annual influenza immunization campaign, or (4) the management of influenza and its potential complications.




Figure 5. Weekly calls to Health Links – Info Santé, Manitoba, 2016-2017

ILI

ILI visits to sentinel physicians

Manitoba participates in *FluWatch*, the Canada’s national surveillance system co-ordinated by Public Health Agency of Canada (PHAC), which monitors the spread of influenza and ILI on a year-round basis. *FluWatch*

consists of a network of laboratories, hospitals, doctor’s offices and provincial and territorial ministries of health. In 2016–2017, there were 19 sentinel physicians recruited throughout Manitoba reporting to *FluWatch* weekly. E&S receives weekly reports from *FluWatch* which present the ILI rate for Manitoba and for each of the participating sentinel physicians. Note that the reporting sentinel physicians are different by week and their reports may not be representative of ILI activity across the province.




Figure 6. Weekly percentage of patient visits to sentinel physicians due to ILI, Manitoba, 2016–2017

ILI visits to Emergency Rooms

Daily ILI related visits to Emergency Department (ED) at Winnipeg Regional Health Authority (WRHA) are submitted to E&S weekly. ILI cases are defined as patients whose triage chief complaints contain either of these symptoms: weakness, shortness of breath, cough, headache, fever, cardiac/respiratory arrest, sore throat, and upper respiratory tract infection complaints.




Figure 7. Weekly visits due to ILI in Emergency Rooms, WRHA, Manitoba

Antiviral dispensing

The units of antiviral drug, Oseltamivir (Tamiflu®), dispensed since October, 1, 2015 to Manitoba residents during the influenza season are reported to E&S from Drug Programs Information Network (DPIN) on a weekly basis. Only drugs dispensed from community retail pharmacies were included in this report. Antiviral drugs dispensed to in-patients or through nursing stations could not be included due to lack of data.




Figure 8. Weekly units of Oseltamivir dispensed, Manitoba, 2016-2017

Antiviral Resistance

Influenza and Respiratory Viruses Section of National Microbiology Laboratory (NML) undertakes enhanced surveillance, investigations, and research on influenza and other respiratory pathogens. A random sample of positive influenza specimens isolated by culture is referred from each provincial laboratory to NML for strain characterization and antiviral resistance testing. The aggregate level information is then shared with provinces and territories on a weekly basis.

Table 1. Antiviral resistance of isolates by influenza type and subtype since September 1, 2016 in Canada and Manitoba, 2016-2017

		Oseltamivir		Zanamivir		Amantadine	
		# Resistant	# Sensitive	# Resistant	# Sensitive	# Resistant	# Sensitive
Canada	A(H3N2)	0	196	0	196	109	0
	A(H1N1)	0	9	0	9	6	0
	B	0	18	0	18	N/A	N/A
Manitoba	A(H3N2)	0	5	0	5	5	0
	A(H1N1)	0	0	0	0	0	0
	B	0	2	0	2	N/A	N/A

Immunization

As per World Health Organization (WHO), all seasonal quadrivalent influenza vaccines for 2016-2017 in the northern hemisphere contain:

- A/Hong Kong/4801/2014 (H3N2)-like virus
- A/California/7/2009 (H1N1)pdm09-like virus
- B/Brisbane/60/2008-like virus
- B/Phuket/3073/2013-like virus

For the 2016-2017 influenza season, MHSAL has been allotted the quadrivalent inactivated vaccines (QIV), Fluzone® Quadrivalent (Sanofi Pasteur) and FluLaval Tetra® (GlaxoSmithKline), and quadrivalent live attenuated influenza vaccine (QLAIV) FluMist® Quadrivalent (AstraZeneca), as part of the province’s Publicly-Funded Seasonal Influenza Immunization Program.

Circulating Strain

NML antigenically characterizes influenza viruses received from Canadian laboratories year-round. In Manitoba, a random sample of positive influenza specimens isolated by culture is referred from CPL to NML for strain characterization.

Since September 1, 2016, NML has characterized 216 influenza A and B viruses.

1. 188 influenza A(H3N2) viruses:
 - 68 influenza A(H3N2) viruses characterized were antigenically similar to A/Hong Kong/4801/2014, the influenza A(H3N2) component of the 2016-17 Northern Hemisphere influenza vaccine. Of those viruses characterized, 53 belonged to genetic group 3C.2a and 15 belonged to genetic group 3C.3a.
 - 120 influenza A(H3N2) viruses did not grow to sufficient hemagglutination titers for antigenic characterization by hemagglutination inhibition (HI) assays. Therefore, genetic characterization was performed. Sequence analysis of the hemagglutinin (HA) gene of these viruses showed that 120 H3N2 viruses belonged to genetic group 3C.2a. The vaccine strain, A/Hong Kong/4801/2014(H3N2)-like virus, also belongs to genetic group 3C.2a.
2. 10 influenza A(H1N1) viruses:
 - 10 influenza A(H1N1) viruses characterized were antigenically similar to A/California/7/2009, the influenza A(H1N1) component in the vaccine.
3. 18 influenza B viruses:
 - 9 influenza B viruses antigenically similar to B/Brisbane/60/2008-like (Victoria lineage), the influenza B component in the vaccine.
 - 9 influenza B viruses were characterized as B/Phuket/3073/2013 (Yamagata lineage), the influenza B component in the quadrivalent vaccine.

Table 2. Influenza Strain Characterization reported by NML since September 1, 2016, Canada, 2016–2017

Strain	Number of viruses	
	Canada	Manitoba
A/Hong Kong/4801/2014 (H3N2)-like	68	4
A/California/7/2009(H1N1)-like	10	0
B/Brisbane/60/2008-like	9	0
B/Phuket/3073/2013-like	9	2

Abbreviations

- CPL = Cadham Provincial Laboratory
- E&S = Epidemiology and Surveillance
- ICU = Intensive Care Unit
- ILI = Influenza-Like-Illness
- LTCF = Long Term Care Facility
- MHSAL= Manitoba Health, Seniors and Active Living
- NML = National Microbiology Laboratory
- PHAC = Public Health Agency of Canada
- RHA = Regional Health Authority
- RSV = Respiratory Syncytial Virus
- WRHA = Winnipeg Regional Health Authority

Explanatory Notes and Definitions

Cumulative data

Cumulative data include updates to previous weeks; due to reporting delays or amendments, the sum of weekly report totals may not add up to cumulative totals.

Data extraction date

Manitoba-specific information contained within this update is based on data confirmed in SIS databases at 11:00 am on the date of data extraction.

Epidemiology week

Time trends in this report were analyzed by [epidemiology week](#), a schedule used by the national FluWatch program coordinated by the Public Health Agency of Canada (PHAC).

Incidence rate

Incidence rate measures the frequency with which influenza occurs in a region. It is calculated as the total number of new cases this influenza season multiplied by 100,000 and divided by the total population in each region. Regional populations are based on the Manitoba Health Population Report 2015.

ILI in the general population

Acute onset of respiratory illness with fever and cough and with one or more of the following – sore throat, arthralgia, myalgia, or prostration, which is likely due to influenza. In children under 5, gastrointestinal symptoms may also be present. In patients under 5 or 65 and older, fever may not be prominent.

ILI outbreaks

Schools: Greater than 10% absenteeism (or absenteeism that is higher (e.g. >5-10%) than expected level as determined by school or public health authority) which is likely due to ILI.

Hospitals and residential institutions: Two or more cases of ILI within a seven-day period.

Other settings: Two or more cases of ILI within a seven-day period, including at least one laboratory confirmed case; i.e. workplace, closed communities.

Specimen collection date

The date the laboratory specimen was taken is used to assign cases to the appropriate week in this report. Occasionally, if the specimen collection date is not available, the laboratory report date will be used.

Epidemiology and Surveillance
Manitoba Health, Seniors and Active Living
flusurveillance@gov.mb.ca
(204)786-7335

For other Epidemiology and Surveillance reports, please view the Manitoba Health internet website:
<http://www.gov.mb.ca/health/publichealth/surveillance/index.html>

For national surveillance data, refer to:
<http://www.phac-aspc.gc.ca/fluwatch/index-eng.php>